

EQUIVARIANT CELLULAR HOMOLOGY AND ITS APPLICATIONS

BORIS CHORNY

ABSTRACT. In this work we develop a cellular equivariant homology functor and apply it to prove an equivariant Euler-Poincaré formula and an equivariant Lefschetz theorem.

1. INTRODUCTION

Let D be an arbitrary small topologically enriched category. In this paper we develop a D -CW-homology functor which allows for easy computation of the *ordinary* D -equivariant homology defined by E. Dror Farjoun in [1]. Our approach is a generalization of the G -CW-(co)homology functor constructed by S.J. Willson in [13] for the case of G being compact Lie group.

Then we apply the D -CW-homology functor to obtain:

(i) Equivariant Euler-Poincaré formula:

$$(1) \quad \chi^D(\underline{X}) = \sum_{n=0}^{\infty} (-1)^n \widetilde{rk}_{\text{HS}}(H_n^D(\underline{X}; \mathcal{I}))$$

This formula establishes a connection between the equivariant homology and an equivariant Euler characteristic; $\widetilde{rk}_{\text{HS}}(\cdot)$ is a slight modification of Hattori-Stallings rank (originally defined in [8],[11]).

(ii) Equivariant Lefschetz theorem: Let $\underline{X}$ be a triangulated D -space, $f : \underline{X} \rightarrow \underline{X}$ an equivariant map. If the equivariant Lefschetz number

$$(2) \quad \Lambda_D(f) = \sum_{n=0}^{\infty} (-1)^n \widetilde{tr}_{\text{HS}}(H_n^D(f; \mathcal{I}))$$

is not equal to zero, then there are f -invariant orbits in $\underline{X}$, moreover the orbit types of the invariant orbits may be recovered from $\Lambda_D(f)$.

Acknowledgments. I would like to thank E.Dror Farjoun, V.Halperin, A.Libman, Sh. Rosset for many stimulating conversations and helpful ideas.

2. PRELIMINARIES

2.1. D -spaces. Let $\mathcal{Top}$ denote the category of the compactly generated Hausdorff topological spaces. Fix an arbitrary small category D enriched over $\mathcal{Top}$. We work in the category $\mathcal{Top}^D$ of functors from D to $\mathcal{Top}$. The objects of this category

Date: August 30, 2001.

1991 *Mathematics Subject Classification.* Primary 55N91; Secondary 55P91, 57S99.

The author acknowledges the support of Sonderforschungsbereich (SFB) 478 of the University of Münster and Edmund Landau Center for Research in Mathematical Analysis.

are called topological diagrams or just D -spaces. The arrows in $\mathcal{Top}^D$ are natural transformations of functors or *equivariant maps*.

2.2. D -homotopy. An equivariant homotopy between two D -maps $f, g : \underline{X} \rightarrow \underline{Y}$, where $\underline{X}, \underline{Y}$ are D -diagrams, is a D -map $H : \underline{X} \times I \rightarrow \underline{Y}$, where I denotes the constant D -space $I(d) = [0, 1]$. A *homotopy equivalence* $f : \underline{X} \rightarrow \underline{Y}$ is a map with a (two sided) D -homotopy inverse.

2.3. D -orbits. We recall now the central concept of the D -homotopy theory (introduced in [1],[3]) – that of D -orbit. A D -orbit is a D -space $T : D \rightarrow \mathcal{Top}$, such that $\text{colim}_D T = \{*\}$. A **free** D -orbit generated in $d \in \text{ob}(D)$ is $\mathcal{Top}^D \ni F^d = \text{hom}_D(d, \cdot)$, i.e. $F^d(d') = \text{hom}_D(d, d')$ and $F^d(d' \rightarrow d'')$ is given by the composition. Clearly F^d is a D -orbit. A D -space $\underline{X}$ is called free iff for any $s \in \text{colim}_D \underline{X}$ the full orbit T_s lying over s is free.

2.4. D -CW-complexes. A D -cell is a D -space of the form $T \times e^n$, where T is a D -orbit and e^n is the standard n -cell. An *attaching map* of this D -cell to some D -space $\underline{X}$ is a map $\phi : T \times \partial e^n \rightarrow \underline{X}$.

A (relative) D -CW-complex $(\underline{X}, \underline{X}_{-1})$ is a D -space $\underline{X}$ together with a filtration $\underline{X}_{-1} \subset \underline{X}_0 \subset \dots \subset \underline{X}_n \subset \underline{X}_{n+1} \subset \dots \subset \underline{X} = \text{colim}_n \underline{X}_n$, such that $\underline{X}_{n+1}$ is obtained from $\underline{X}_n$ by attaching a set of n -dimensional D -cells. Namely one has a push-out diagram of D -spaces:

$$\begin{array}{ccc} \coprod_i (T_i \times \partial e^n) & \xrightarrow{\phi} & \underline{X}_{n-1} \\ \downarrow & & \downarrow \\ \coprod_i (T_i \times e^n) & \xrightarrow{\Phi} & \underline{X}_n \end{array}$$

If $\underline{X}_{-1} = \emptyset$ we call the D -CW-complex *absolute*.

Let $\underline{X}$ be a D -CW-complex. A D -subspace $\underline{Y} \subset \underline{X}$ is called the *cellular subspace* if $\underline{Y}$ has a D -CW-structure such that each cell of $\underline{Y}$ is also a cell of $\underline{X}$.

2.5. The category of orbits. The *category of orbits* $\mathcal{O}$ is a full topological subcategory of $\mathcal{Top}^D$ generated by all D -orbits.

Usually $\mathcal{O}$ is not a small category. For example for $D = J = (\bullet \rightarrow \bullet)$, then $\mathcal{O} \cong \mathcal{Top}$. A model category has been constructed for the D -spaces of arbitrary orbit type in [2]. We will be interested in the diagrams which are homotopy equivalent to the finite D -CW-complexes, i.e. only finite number of orbit types appear in such diagram. We collect those orbits into the a full subcategory $\mathcal{O}'$ of $\mathcal{O}$ with a finite amount of objects.

2.6. Orbit point $(\cdot)^\mathcal{O}$ and realization $|\cdot|_D$ functors. Suppose $\mathcal{O}$ is a *small* category of D -orbits. An orbit point functor $(\cdot)^\mathcal{O} : \mathcal{Top}^D \rightarrow \mathcal{Top}^{\mathcal{O}^{\text{op}}}$ is a generalization to the diagram case of Bredon's fixed point functor. For any D -space $\underline{X}$ (usually of type $\mathcal{O}$) $(\underline{X})^\mathcal{O}$ is a $\mathcal{O}^{\text{op}}$ diagram such that $(\underline{X})^\mathcal{O}(T) = \text{hom}_D(T, \underline{X})$ for all $T \in \text{ob}(\mathcal{O})$ and the arrows of the diagram are induced by composition with the maps between orbits.

If $f : \underline{X} \rightarrow \underline{Y}$ is an equivariant map between two D -spaces, then there exist an $\mathcal{O}^{\text{op}}$ -equivariant map $f^\mathcal{O} : (\underline{X})^\mathcal{O} \rightarrow (\underline{Y})^\mathcal{O}$, which is obtained from f by composition:

$$(\underline{X})^\mathcal{O}(T) = \text{hom}_D(T, \underline{X}) \ni g \xrightarrow{f^\mathcal{O}} f \circ g \in \text{hom}_D(T, \underline{Y}) = (\underline{Y})^\mathcal{O}(T)$$

The fundamental property of $(\cdot)^\mathcal{O}$ functor is that for any D -space $\underline{X}$ the $\mathcal{O}^\text{op}$ -space $(\underline{X})^\mathcal{O}$ is $\mathcal{O}^\text{op}$ -free [1, 3.7].

Example 2.1. Consider a free D -space $\underline{X}$. And let the orbit category $\mathcal{O}$ consists of all the free orbits. Then $\mathcal{O}$ is isomorphic to D^op as a category and $(\underline{X})^\mathcal{O} \cong \underline{X}$ (Yoneda's lemma).

Another easy case occurs then $\underline{X}$ is a D -CW-space. We shall discuss it in the next section.

There exist a left adjoint to $(\cdot)^\mathcal{O}$. It is called *realization functor* $|\cdot|_D$, since it takes an $\mathcal{O}^\text{op}$ -space and produce a D -space with the prescribed orbit point data (up to local weak equivalence). Realization functor in the group case has been constructed by A.D.Elmendorf in [7] and has been generalized to the arbitrary diagram case by W.Dwyer and D.Kan in [6]. Compare also [1].

2.7. Equivariant Euler characteristic. Let $\underline{X}$ be a finite D -CW-complex, then $\underline{X}$ is of type $\mathcal{O}$ for some category of orbits with finite amount of objects. We define the Equivariant Euler characteristic to be the Universal Additive Invariant [10, I.5] $(U(D), \chi^D)$. Or, equivalently, we say that $\chi^D(\underline{X}) \in U(D)$ is equal to the alternating sum of the orbit types over the dimensions of the cells of $\underline{X}$ in the free abelian group $U(D) = \bigoplus_{T \in \text{Iso}(h\mathcal{O})} \mathbb{Z}$ generated by the homotopy types of orbits.

3. EQUIVARIANT CELLULAR HOMOLOGY

3.1. $\mathcal{O}^\text{op}$ -CW structure on the orbit point space of a D -CW-complex. Let $\mathcal{C}$ denote a subcategory of $\text{Top}^{\mathcal{O}^\text{op}}$ which is obtained as the image of Top^D under the functor $(\cdot)^\mathcal{O}$. Recall that there is the inclusion of the categories $\iota : D \hookrightarrow \mathcal{O}^\text{op}$, where $\iota(d) = F^d$, for each $d \in \text{ob}(D)$. Hence there is a functor $\text{Res} : \text{Top}^{\mathcal{O}^\text{op}} \rightarrow \text{Top}^D$. By abuse of notation we denote by Res also $\text{Res}|_{\mathcal{C}}$.

Lemma 3.1. *The functor $(\cdot)^\mathcal{O}$ is fully faithful.*

Proof. The faithfulness is clear. We have to show only that for any map $f_0 : \underline{X}^\mathcal{O} \rightarrow \underline{Y}^\mathcal{O}$ there exists a map $f : \underline{X} \rightarrow \underline{Y}$ s.t. $f_0 = f^\mathcal{O}$. Take $f = \text{Res}(f_0)$, then if $\underline{X}$ and $\underline{Y}$ were orbits the result follows from the bijective correspondence induced by the Yoneda's lemma: $\text{hom}_D(\underline{X}, \underline{Y}) = \text{hom}_{\mathcal{O}^\text{op}}(\underline{X}^\mathcal{O}, \underline{Y}^\mathcal{O})$. The general claim will follow from the comparison of f_0 and $f^\mathcal{O}$ orbitwise, i.e. by their action on each full orbit. Fortunately the functor $(\cdot)^\mathcal{O}$, being right adjoint, commutes with taking full orbit (pullback). $\square$

Lemma 3.2. *The pair of functors $\text{Res}(\cdot) : \mathcal{C} \leftrightarrow \text{Top}^D : (\cdot)^\mathcal{O}$ induce the equivalence of the categories $\mathcal{C}$ and Top^D .*

Proof. We need to construct the natural isomorphisms of the functors $\text{id}_{\text{Top}^D} \cong \text{Res}((\cdot)^\mathcal{O})$ and $\text{id}_{\mathcal{C}} \cong (\text{Res}(\cdot))^\mathcal{O}$.

Let $\underline{X} \in \text{Top}^D$, then $\text{Res}(\underline{X}^\mathcal{O}) \cong \underline{X}$ because the generalized lemma of Yoneda [9] induces the objectwise homeomorphisms and the equivariance is preserved by the naturality of the Yoneda's isomorphism. But an equivariant map which is the objectwise homeomorphism is an isomorphism of D -spaces, hence the first isomorphism of functors.

Let $\underline{X}^\mathcal{O} \in \mathcal{C}$, then $\text{Res}(\underline{X}^\mathcal{O}) \cong \underline{X}$ by the first homeomorphism, then $(\text{Res}(\underline{X}^\mathcal{O}))^\mathcal{O} \cong \underline{X}^\mathcal{O}$. Hence the second isomorphism. $\square$

Proposition 3.3. *Let $\underline{X}$ be a (pointed) D -CW-space of orbit type $\mathcal{O}$, where $\mathcal{O}$ is a small category of orbits. Consider the $\mathcal{O}^{\text{op}}$ -space $\underline{X}^{\mathcal{O}}$ to be the orbit point space of $\underline{X}$.*

Then $\underline{X}^{\mathcal{O}}$ has $\mathcal{O}^{\text{op}}$ -CW structure which corresponds to the D -CW structure of $\underline{X}$ in the following sense: let $pt_D \subseteq \underline{X}_0 \subseteq \underline{X}_1 \subseteq \cdots \subseteq \underline{X}_n \subseteq \cdots \subseteq \underline{X} = \text{colim}_n \underline{X}_n$ is a D -CW filtration of $\underline{X}$, such that each $\underline{X}_n$ is a push-out:

$$\begin{array}{ccc} \coprod_i T_i \times S^{n-1} & \xrightarrow{\phi} & \underline{X}_{n-1} \\ \downarrow & & \downarrow i_n \\ \coprod_i T_i \times D^n & \xrightarrow{\Phi} & \underline{X}_n \end{array}$$

then there exist a $\mathcal{O}^{\text{op}}$ -CW-filtration: $pt_{\mathcal{O}^{\text{op}}} \subseteq \underline{X}_0^{\mathcal{O}} \subseteq \underline{X}_1^{\mathcal{O}} \subseteq \cdots \subseteq \underline{X}_n^{\mathcal{O}} \subseteq \cdots \subseteq \underline{X}^{\mathcal{O}} = \text{colim}_n \underline{X}_n^{\mathcal{O}}$, such that $\underline{X}_n^{\mathcal{O}} = (\underline{X}_n)^{\mathcal{O}}$, and

$$\begin{array}{ccc} \coprod_i F^{T_i} \times S^{n-1} & \xrightarrow{\phi^{\mathcal{O}}} & \underline{X}_{n-1}^{\mathcal{O}} \\ \downarrow & & \downarrow i_n^{\mathcal{O}} \\ \coprod_i F^{T_i} \times D^n & \xrightarrow{\Phi^{\mathcal{O}}} & \underline{X}_n^{\mathcal{O}} \end{array}$$

is a push-out square.

Proof. We proceed by the induction on the skeleton of $\underline{X}$.

$$\begin{aligned} \underline{X}_0^{\mathcal{O}}(T) &= (\coprod_i T_i)^{\mathcal{O}}(T) = \coprod_i ((T_i)^{\mathcal{O}}(T)) = \coprod_i \text{hom}_D(T, T_i) = \\ &= \coprod_i \text{hom}_{\mathcal{O}^{\text{op}}}(F^T, F^{T_i}) = \coprod_i F^{T_i}(T). \end{aligned}$$

Hence the base of the induction.

Suppose we know the claim for $\underline{X}_n$. Then it follows for $\underline{X}_{n+1}$ since $(\cdot)^{\mathcal{O}}$ is both left and right adjoint, so commutes both with push-outs and products. $\square$

3.2. D -CW-homology functor. The construction of the (co)homology functor in [1, 4.16] depends on the specific D -CW-decomposition of $\underline{X}^{\mathcal{O}}$. We apply this construction to the cellular structure of $\underline{X}^{\mathcal{O}}$, which was constructed in 3.3 and obtain the required D -CW-homology functor.

3.3. Isotropy ring $\mathcal{I}$. In [13] a universal coefficient system for the G -equivariant homology have been developed (where G is a compact Lie group). Let us generalize this approach to the coefficient systems for the classical D -homology theory. Suppose $\mathcal{O}'$ is a small, full subcategory of the orbit category $\mathcal{O}$. Let $\underline{X}$ be a D -space of orbit type $\mathcal{O}'$. Then a coefficient system for the ordinary (co)homology is a homotopy (co)functor $M : \mathcal{O}' \rightarrow (R - \text{mod})$.

Definition 3.4. Let R be a commutative ring. An *isotropy ring* $\mathcal{I} = I_D^{R, \mathcal{O}'}$ is generated by $\text{mor}(h\mathcal{O}')$ as a free $R - \text{mod}$. Define the multiplication on the generators by

$$fg = \begin{cases} f \circ g, & \text{if } \text{codom}(g) = \text{dom}(f) \\ 0, & \text{otherwise} \end{cases}$$

and extend the definition to the rest of the elements of $\mathcal{I}$ by linearity.

Proposition 3.5. *The category $\mathcal{M}$ of the left $\mathcal{I}$ -modules which satisfy:*

$$(3) \quad \forall M \in \text{ob}(\mathcal{M}), M = \bigoplus_{T \in \text{ob}(h\mathcal{O}')} 1_T M$$

(where $\{1_T M\}_{T \in \text{ob}(h\mathcal{O}')}$ are left R -modules) and the category of $R(h\mathcal{O}')$ -mod of functors from $h\mathcal{O}'$ to the category of left R -modules are equivalent.

Proof. Let us define a pair of functors which induce the required equivalence:

$$\zeta : \mathcal{M} \rightleftharpoons R(h\mathcal{O}')\text{-mod} : \xi.$$

Let $M \in \text{ob}(\mathcal{M})$, $T \in \text{ob}(h\mathcal{O}')$, then define

$$\zeta M(T) = 1_T M.$$

If $\text{mor}(h\mathcal{O}') \ni f : T_1 \rightarrow T_2$, then define

$$\zeta M(f)(1_{T_1} m) = f 1_{T_1} m = (1_{T_2} f) 1_{T_1} m \in 1_{T_2} M.$$

Obviously the morphisms of the left $\mathcal{I}$ modules correspond to the natural transformations of the functors.

Given a $R(h\mathcal{O}')$ -module N , then

$$\xi N = \bigoplus_{T \in \text{ob}(h\mathcal{O}')} N(T), \text{ as a left } R\text{-module.}$$

Define the left $\mathcal{I}$ -module structure on ξN by $f(\dots, n, \dots) = (\dots, f n, \dots)$, where

$$N(\text{codom}(f)) \ni f n = \begin{cases} f(n), & \text{if } n \in N(\text{dom}(f)) \\ 0, & \text{otherwise} \end{cases}$$

Now it is clear that the defined functors provide the equivalence of the categories. $\square$

Remark 3.6. The ring $\mathcal{I}$ considered as a left $\mathcal{I}$ -module is an object of $\mathcal{M}$, because $\mathcal{I} \cong \bigoplus_{T \in \text{ob}(h\mathcal{O}')} 1_T \mathcal{I}$ (as left R -modules) by the construction. But it also carries an obvious structure of the right $\mathcal{I}$ module, so the $\zeta \mathcal{I}(T)$.

Remark 3.7. If $\text{ob}(h\mathcal{O}')$ is finite then the ring $\mathcal{I}$ has a two-sided identity element $1 = \sum_{T \in \text{ob}(h\mathcal{O}')} 1_T$ together with its decomposition into the sum of the orthogonal idempotents and the condition (3) is redundant.

Definition 3.8. The augmentation $\varphi : \mathcal{I} \rightarrow \bigoplus_{T \in \text{Iso}(\text{ob}(h\mathcal{O}'))} R$ is defined for any

$$\mathcal{I} \ni g = \sum_{T \in \text{ob}(h\mathcal{O}')} \sum_{f \in \text{mor}(T, T)} r_f f + \sum_{h \in \text{mor}(T_1, T_2), T_1 \neq T_2} s_h h$$

(only a finite number of $r_f, s_h \in R$ is non equal to zero) to be

$$\varphi(g) = (\dots, \sum_{f \in \text{mor}(T, T)} r_f, \dots) \in \bigoplus_{T \in \text{Iso}(\text{ob}(h\mathcal{O}'))} R$$

Remark 3.9. The idempotents in $\mathcal{I}$ which correspond to the D -homotopy equivalent orbits are identified under φ . Apparently, φ is an epimorphism of rings. Consider the abelinization functor $Ab : (Rings) \rightarrow \mathcal{A}b$ which corresponds to a ring its additive group divided by the commutator subgroup. Then $Ab(\varphi) : Ab(\mathcal{I}) \rightarrow \bigoplus_{T \in \text{Iso}(\text{ob}(h\mathcal{O}'))} R$. The last map will be used to obtain a generalization of the Euler-Poincaré formula.

4. APPLICATIONS

Let $\underline{X}$ be a finite D -CW-complex of type $\mathcal{O}'$ for some orbit category $\mathcal{O}'$ with $\text{obj}(\mathcal{O}')$ a finite set.

4.1. Equivariant Euler-Poincaré formula. We remind that the equivariant Euler characteristic lies in the abelian group $U(D) \cong \bigoplus_{Iso(\text{obj}(h\mathcal{O}'))} \mathbb{Z}$, so in order to apply Hattori–Stallings machinery we need to choose a coefficient system for the equivariant homology such that the resulting chain complex and homology groups will be endowed with the module structure over some ring S which allows an epimorphism $\varepsilon : Ab(S) \rightarrow U(D)$.

Our choice of the coefficient system for the equivariant homology will be the isotropy ring $\mathcal{I} = I_D^{\mathbb{Z}, \mathcal{O}'}$ taken over itself as a left module.

Lemma 4.1. *Let $\underline{X}$ be a finite D -CW-complex. Suppose $\underline{X}$ has n_q q -dimensional cells and $t_1 + \dots + t_s = n_q$, t_i is the number of q -dimensional cells of the same homotopy type $T_i \in Iso(\text{obj}(h\mathcal{O}'))$. Then $\mathcal{C}_q(\underline{X}) \otimes_{\mathcal{O}'} \zeta \mathcal{I} \cong \zeta \mathcal{I}(T_1)^{t_1} \oplus \dots \oplus \zeta \mathcal{I}(T_s)^{t_s}$ as a left $\mathbb{Z}$ -module.*

Proof. Let $t_i = r_{i1} + \dots + r_{ik}$, where r_{ij} is the number of q -dimensional cells of type $T_{ij} \in \text{ob}(\mathcal{O})$ of homotopy type T_i . By the construction of the equivariant homology $\mathcal{C}_q(\underline{X}) = \bigoplus_{i=1}^s (\bigoplus_{j=1}^k \mathbb{Z}(\text{hom}_{\mathcal{O}'}(?, T_{ij})^{r_{ij}}))$. The dual Yoneda isomorphism [9, p.74] implies:

$$\mathcal{C}_q(\underline{X}) \otimes_{\mathcal{O}'} \zeta \mathcal{I} \cong \bigoplus_{i=1}^s (\bigoplus_{j=1}^k \zeta \mathcal{I}(T_{ij})^{r_{ij}}) \cong \bigoplus_{i=1}^s \bigoplus_{j=1}^k (1_{T_{ij}} \mathcal{I})^{r_{ij}},$$

If T_{ij_1} is isomorphic to T_{ij_2} in $h\mathcal{O}'$ then there is an obvious isomorphism of the left $\mathbb{Z}$ -modules and right $\mathcal{I}$ -modules $1_{T_{ij_1}} \mathcal{I} \cong 1_{T_{ij_2}} \mathcal{I}$. Let us choose a representative T_i of each isomorphism class of objects in $h\mathcal{O}'$, then

$$\mathcal{C}_q(\underline{X}) \otimes_{\mathcal{O}'} \zeta \mathcal{I} \cong \bigoplus_{i=1}^s (1_{T_i} \mathcal{I})^{(\sum_{j=1}^k r_{ij})} \cong \bigoplus_{i=1}^s (1_{T_i} \mathcal{I})^{t_i} \cong \bigoplus_{i=1}^s (\zeta \mathcal{I}(T_i))^{t_i}$$

□

Because of 3.6 the equivariant chain complex $\{\mathcal{C}_q(\underline{X}) \otimes_{\mathcal{O}'} \zeta \mathcal{I}\}_{q=0}^{\dim \underline{X}}$ is a complex of *projective* right $\mathcal{I}$ -modules and the equivariant homology is endowed with the right $\mathcal{I}$ -module structure.

Notation: $\chi_{\text{HS}}(\cdot)$ means Euler characteristic of a $\mathcal{I}$ differential complex with respect to $rk_{\text{HS}}(\cdot)$.


Proposition 4.2. *Let $K_* = \mathcal{C}_*(\underline{X}) \otimes_{\mathcal{O}'} \zeta \mathcal{I}$ be a right $\mathcal{I}$ -complex, then $\chi^D(\underline{X}) = Ab(\varphi)(\chi_{\text{HS}}(K_*))$ whenever left side is defined.*

Proof. It is easy to see that $rk_{\text{HS}}(1_T \mathcal{I}) = 1_T \in Ab(\mathcal{I})$. Lemma 4.1 together with 3.9 completes the proof. □

Now we combine 4.2 with the additivity properties of the Hattori–Stallings rank and obtain the following

Theorem 4.3. $\chi^D(\underline{X}) = Ab(\varphi)(\sum_{n=0}^{\infty} (-1)^n rk_{\text{HS}} H_n^D(\underline{X}; \zeta \mathcal{I}))$, whenever the left side is defined.

Example 4.4. Consider the J -diagram:



$\underline{Z}$ has two 0-cells of type $T_2 = [\downarrow]$ and one 1-cell of type $T_3 = [\ddownarrow]$, hence $\chi^J(\underline{Z}) = 2[\downarrow] - [\ddownarrow]$.

The category $\mathcal{O}'$ of orbits contains two objects: T_2, T_3 . The cellular chain complex tensored with the coefficients $\mathcal{I} = \mathcal{I}_J^{\mathbb{Z}, \{T_2, T_3\}}$ becomes:

$$\cdots \rightarrow 0 \rightarrow 1_{T_3}\mathcal{I} \xrightarrow{\partial_1} (1_{T_2}\mathcal{I})^2$$

and $\partial_1 = 0$ from the orbit type considerations.

$U(J) = \mathbb{Z} \oplus \mathbb{Z}$ in that case. And $H_0^J(\underline{Z}, \mathcal{I}) = (1_{T_2}\mathcal{I})^2$, $H_1^J(\underline{Z}, \mathcal{I}) = 1_{T_3}\mathcal{I}$ are right $\mathcal{I}$ -modules. Hence, $\chi^J(\underline{Z}) = (2, 0) - (0, 1) = (2, -1)$.

Let us, for comparison, calculate the J -equivariant homology of $\underline{Z}$ with $\mathbb{Z}^{\mathcal{O}'}$ coefficients: $H_i^J(\underline{Z}, \mathbb{Z}^{\mathcal{O}'}) = H_i(\text{colim}_J \underline{Z}, \mathbb{Z})$ (see [1, 5.2]). Then $\text{colim}_J \underline{Z} = I = [0, 1]$ and

$$H_i^J(\underline{Z}, \mathbb{Z}^{\mathcal{O}'}) = \begin{cases} \mathbb{Z}, & i = 0 \\ 0, & \text{otherwise} \end{cases}$$

We can see that $\mathbb{Z}^{\mathcal{O}'}$ coefficients are inappropriate to the Euler-Poincaré formula.

4.2. Equivariant Lefschetz theorem. Using cellular equivariant homology functor we are able now to proof a version of the equivariant Lefschetz theorem.

Some result of the Lefschetz type in the equivariant setting may be obtained already by applying the ordinary Lefschetz theorem: consider an equivariant map $f : \underline{X} \rightarrow \underline{X}$, where $\underline{X}$ is a diagram over small category D , then if the Lefschetz number $\Lambda(\text{colim}_D \underline{X}) \neq 0$ there are f -invariant D -orbits in $\underline{X}$. However the advantage of using the equivariant homology and equivariant Lefschetz number $\Lambda_D(\underline{X}) \in U(D)$ is that we obtain the specific information about orbit type of the invariant orbit.

First we give a technical

Definition 4.5. A D -CW-complex $\underline{X}$ will be called the triangulated D -space if the natural CW -structure of $\text{colim } \underline{X}$ also triangulates $\text{colim } \underline{X}$.

The following lemma will be used in the proof of the equivariant Lefschetz theorem.

Lemma 4.6. Let $\underline{X}$ be a triangulated diagram, then for any refinement Y of the triangulation of $\text{colim } \underline{X}$, there exists a D -CW-complex $\underline{X}'$, such that $\underline{X}'$ is D -homeomorphic to $\underline{X}$ and $\text{colim } \underline{X}' = Y$ (as the triangulated spaces). $\underline{X}'$ will be called the refinement of $\underline{X}$.

Proof. Consider a new simplex Δ in the triangulation of Y . It lies in some old simplex of $\text{colim } \underline{X} : \Delta \in \Delta'$. Then consider the pull-back:

$$\lim \left(\begin{array}{ccc} & \underline{X} & \\ & \downarrow & \\ \Delta & \hookrightarrow & \text{colim}_D \underline{X} \end{array} \right) = T \times \Delta,$$

where T is the orbit which lies over Δ' .

We've obtained the cell of the new D -CW-complex $\underline{X}'$. Continuing in the same way for the rest of the simplices of Y completes the construction of $\underline{X}'$. Hence D -CW-complex $\underline{X}'$ has the same underlying topological diagram as $\underline{X}$, therefore they are D -homeomorphic. $\square$

Definition 4.7. Let $f : \underline{X} \rightarrow \underline{X}$ be a map of the finite triangulated D -space $\underline{X}$ of orbit type $\mathcal{O}'$, where $\mathcal{O}'$ is an orbit category with the finite number n of objects. Let $\mathcal{I} = \mathcal{I}_D^{\mathbb{Z}, \mathcal{O}'}$. Then the equivariant Lefschetz number of f :

$$U(D) \ni (\lambda_1, \dots, \lambda_n) = \Lambda_D(f) = Ab(\varphi) \left(\sum_{k=0}^{\infty} (-1)^k \text{tr}_{\text{HS}}(H_k(f; \mathcal{I})) \right)$$

Theorem 4.8. Let $\underline{X}$ be a finite triangulated diagram over D . $f : \underline{X} \rightarrow \underline{X}$ be a D -map. $\Lambda_D(f) = (\lambda_1, \dots, \lambda_n) \in U(D)$ - Lefschetz number of f . Then if there is no f -invariant orbit of type T_m , $\lambda_m = 0$.

Proof. A simplex in $\text{colim } \underline{X}$ will be called of type T if the overlaying orbit is of type T in $\underline{X}$. Then the condition that there are no invariant orbits of type T_m is equivalent to the condition that there are no fixed points in the simplices of type T_m .

Since $\underline{X}$ is a finite triangulated diagram, $\text{colim } \underline{X}$ is a finite triangulated space, hence it is a compact metric space. If there are no fixed points of type T_m , then there exists a refinement Y of the triangulation such that if Δ is a simplex of type T_m in Y , $\Delta \cap (\text{colim } f)(\Delta) = \emptyset$.

Consider the refinement $\underline{X}'$ of $\underline{X}$, which exists by lemma 4.6. Since $\underline{X}' \cong \underline{X}$, $H_*^D(\underline{X}'; \mathcal{I}) = H_*^D(\underline{X}; \mathcal{I})$, $\Lambda(f') = \Lambda(f)$, where $f' : \underline{X}' \rightarrow \underline{X}'$ is equal to f , $\lambda'_m = \lambda_m$. Therefore, it is enough to show that $\lambda'_m = 0$.

Now,

$$\Lambda_D(f') = Ab(\varphi) \left(\sum_{k=0}^{\infty} (-1)^k \text{tr}_{\text{HS}}(H_k(f'; \mathcal{I})) \right) = Ab(\varphi) \left(\sum_{k=0}^{\infty} (-1)^k \text{tr}_{\text{HS}}(\mathcal{C}_k(f'; \mathcal{I})) \right),$$

where $\mathcal{C}_k(f'; \mathcal{I})$ is the map induced by f on the chains $\mathcal{C}_k(\underline{X}; \mathcal{I}) = \mathcal{C}_k(\underline{X}) \otimes_{\mathcal{O}'} \mathcal{I} = (1_{T_1} \mathcal{I})^{t_1} \oplus \dots \oplus (1_{T_n} \mathcal{I})^{t_n}$ as $\mathcal{I}$ -module. Because of the property: $\Delta \cap (\text{colim } f)(\Delta) = \emptyset$ for *any* simplex Δ of type T_m , the induced map on $\mathcal{C}_k(\underline{X}; \mathcal{I})$ will take the generator 1_{T_m} corresponding to Δ outside the submodule $1_{T_m} \mathcal{I}$, that it generates. Then the m -th entry of $Ab(\varphi) \left(\sum_{k=0}^{\infty} (-1)^k \text{tr}_{\text{HS}}(\mathcal{C}_k(f'; \mathcal{I})) \right)$ will be zero. This is true for all k , hence $\lambda_m = 0$. $\square$

REFERENCES

- [1] E. Dror Farjoun, *Homotopy and homology of diagrams of spaces*, Proceedings of Conference on Algebraic Topology, Seattle 1985. Lecture Notes in Math.1286, 93–134, Springer-Verlag 1987.

- [2] E. Dror Farjoun, *Homotopy theories for diagrams of spaces*, Proc. Amer. Math. Soc. 101, 181–189, 1987.
- [3] E. Dror Farjoun, A. Zabrodsky, *Homotopy equivalence between diagrams of spaces*, J. Pure Appl. Alg. 41, 169–182, 1986.
- [4] T. tom Dieck, *Transformation groups*, de Gruyter Studies in Mathematics 8, Walter de Gruyter, 1987
- [5] A. Dold, *Lectures on Algebraic topology*, Springer-Verlag, 1972
- [6] W.G. Dwyer, D.M. Kan, *Singular functors and realization functors*, Proc. Kon. Akad. van Wetensch. A87 = Ind.Math. 46, 139–146, 1984.
- [7] A.D. Elmendorf, *Systems of fixed point sets*, Trans. Amer. Math. Soc. 227, 275–284, 1983.
- [8] A. Hattori, *Rank element of a projective module*, Nagoya Math. J. 25, 113–120, 1965.
- [9] G.M. Kelly, *Basic concepts of enriched category theory*, London Math. Soc., Lecture Note Series 64, Cambridge University Press, 1982.
- [10] W. Lück, *Transformation groups and algebraic K-theory*, Lecture Notes in Math. 1408, Springer-Verlag, 1989.
- [11] J.R. Stallings, *Centerless groups—an algebraic formulation of Gottlieb’s theorem*, Topology 4, 129–134, 1965.
- [12] Ch. Watts, *A homology theory for small categories*, Proceedings of the Conference in Categorical Algebra, LaJolla, Springer-Verlag, 1965.
- [13] S.J. Willson, *Equivariant homology theories on G-complexes*, Trans. Amer. Math. Soc. 212, 155–171, 1975.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 91904, ISRAEL

E-mail address: `chorny@math.huji.ac.il`